

# Livestock News

May 2021

## IMPORTANT INFORMATION

### 2021 Regional Chicken Project



The Regional Chicken Project provides an opportunity for youth in both 4-H and FFA to learn how to raise and show chickens. They have the option of raising laying hens or broilers, and complete a project record book documenting their accomplishments and knowledge gained during the project.

Youth had to attend trainings on how to properly complete their project record book and properly show their chickens. The show is the culmination of the project, with youth competing in showmanship classes where they have to demonstrate their knowledge of their bird and ability to properly handle their bird. The chickens are also shown in their respective breed classes and

judged against other birds at the show. The winners of the project record book competition are announced at the show as well.

This year, two youth from Robeson County participated, Jase Chavis and Maggie Love. Jase competed in the Cloverbud division for showmanship and project record books. Maggie competed in the senior division for showmanship and project record books. Jase and Maggie both showed a Barred Rock and Buff Orpington in the showmanship and breed classes. Cloverbuds are non-competitive. Maggie placed fifth in senior showmanship and first in project record books.



### Hay Directory

North Carolina Department of Agriculture's Hay Alert lists people selling hay or looking for hay to buy. It is free to list your hay. To access the directory, visit <http://www.ncagr.gov/HayAlert/>.

### Inside This Issue

- Important Information..... 1
- Animal Waste Management..... 2
- Fly Control Options ..... 3
- Forages for Summer Grazing ..... 4
- Are You Using the Correct Dewormer?..... 5
- Summer Insects and Horses ..... 6
- Youth Livestock Contests - No Critter Required! ..... 7
- USDA Bock Grant for Confined Livestock Operations ..... 8

Robeson County Center  
 O. P. Owens Agriculture Center  
 P. O. Box 2280  
 455 Caton Road  
 Lumberton, NC 28359

Phone - 910-671-3276  
 Fax - 910-671-6278  
 Website - [robeson.ces.ncsu.edu](http://robeson.ces.ncsu.edu)

NC State University and N.C. A&T State University are collectively committed to positive action to secure equal opportunity and prohibit discrimination and harassment regardless of age, color, disability, family and marital status, gender identity, genetic information, national origin, political beliefs, race, religion, sex (including pregnancy), sexual orientation, and veteran status. NC State, N.C. A&T, U.S. Department of Agriculture, and local governments cooperating.

If you are interested in learning more about any information in this newsletter, contact the Extension Center at 910-671-3276 or visit our website at [robeson.ces.ncsu.edu](http://robeson.ces.ncsu.edu). For accommodations for persons with disabilities, contact Cooperative Extension no later than five (5) business days before the event.

Taylor Chavis  
 Extension Agent  
 Agriculture - Livestock



**Animal Waste Management—Anaerobic Digesters**

By: Becky Spearman, Extension Livestock Agent, N.C. Cooperative Extension, Bladen County Center

To check your hours, click [go.ncsu.edu/oichours](http://go.ncsu.edu/oichours), then scroll down and find your name to see number of hours on file, number of hours needed, and date hours are needed by.

**Pumping Considerations**

I talked with some DEQ inspectors about issues they have seen this year. One of the main issues is failure to walk lagoons and failure to walk spray fields before, during, and after waste application. Most of the violations could have been prevented if the Operator in Charge (OIC) would pay attention to what's going on by walking the ENTIRE lagoon and going out in spray fields after spray events.

In addition, if the OIC is not directly pumping and has an employee doing the pumping, the OIC is responsible for inspection of the spray site within 24 hours. As the OIC, you are responsible for the entire farm, even if your employees are doing the pumping, etc. It is the OIC's job to educate their employees and make sure what they should be looking for while pumping. Inspection shall include, but not be limited to, visual observation of application equipment, land application area, subsurface drain outlets, ditches, and drainage ways for any discharge of waste.

Your permit also has that you will inspect the system while pumping at least every 120 minutes during the application of waste. A record of each inspection shall be recorded on IRR forms or other approved forms and includes the date, time, land application area used, and name of the operator for each inspection.

Excessive Ponding means any area of the application field where visible liquid waste is ponded on the surface of the land application site more than four (4) hours following the application of waste. Excessive ponding also means any areas where the ponding of waste has resulted in crop failure.

Record keeping forms. You can use an IRR-1 and IRR-2 form, OR you can use the combined IRR form. All these forms can be downloaded as an Excel spreadsheet, so you can use the formulas to do the math for you. The inspection column includes initials of person inspecting the application site and the time(s) they inspected (at least every 120 minutes/2 hours). Below is an example combined IRR-2 form filled out. You can download many DEQ forms at [go.ncsu.edu/deqforms](http://go.ncsu.edu/deqforms)

**FORM IRR-2**

Lagoon Liquid Irrigation Fields Record  
One Form for Each Field per Crop Cycle

Field Size (wetted acres) = (A)	Zone #	B	Facility Number	9	-	2017
Farm Owner	6		Irrigation Operator			Same
Owner's Address	Becky Spearman		Irrigation Operator's Address			
Owner's Phone #	1234 Hog Way, Hogtown, NC 28337		Operator's Phone #			
	910-876-3623					
<b>From Waste Utilization Plan</b>						
Crop Type	bermuda hay		Recommended PAN Loading (lb/acre)		250	

Lagoon ID	Date (mm/dd/yr)	Start Time	End Time	Irrigation					Waste Analysis PAN* (lb/1000 gal)	PAN Applied (lb/acre) (8) x (9) 1000	Nitrogen Balance** (lb/acre) 250	Weather Code	Inspections (Initials)
				Total Minutes (3) - (2)	# of Sprinklers Operating	Flow Rate (gal/min)	Total Volume (gallons) (6) x (5) x (4)	Volume per Acre (gal/acre) (7) / (A)					
1	4/12/21	2:00 PM	6:00 PM	240	1	230	55200	9200	1.6	14.72	235.28	c	4pm BS, 6pm BS

## Fly Control Options

By: Liz Lahti, Extension Livestock Agent, N.C. Cooperative Extension, Cumberland and Hoke County Centers

The days are getting longer and warmer, which means fly season will soon be upon us! We have learned we will never be able to completely eliminate flies on farms, but we can use a multipronged approach to help keep the population to a tolerable level. There are three main flies of concern and several control options.

The three main flies of concern are horn, face, and stable flies. Horn and stable flies are both blood feeders, with horn flies taking around 30 blood meals per day per animal. Horn flies are usually found on the backs, sides, poll, and bellies of cattle, with stable flies spending most of their time on the cattle's legs. Stable fly bites are very painful and usually cause cattle to stomp their legs when bit. Face flies feed on animal secretions, nectar, and dung liquids, and are best known for spreading the bacteria that causes pinkeye. Face flies will often congregate around eyes and the corner of the cattle's mouths and are an extreme annoyance for the animals.




Photo: John B. Campbell, University of Nebraska-Lincoln

**Figure 2. Comparison of the horn fly to the stable fly, house fly, and face fly.**

There are biological, cultural, and chemical control options for combating fly populations. Predatory wasps are an example of a biological control method but are usually unrealistic for animals that live out on pasture. Removing manure, old hay, and spoiled feed are all cultural control methods that will help reduce where the three flies mentioned above lay their eggs.

Most beef cattle producers will also use chemical control methods. Keep in mind that insecticides are grouped based on how they work against insects. Continued use of products from the same group will lead to the insects developing resistance causing the chemicals to no longer be effective. Because of this, it is important to rotate among the groups. The most commonly used chemical control methods include feeding a larvicide or insect growth regulator (IGR), using fly tags impregnated with insecticide, dust bags/cattle rubs, and sprays.

### Feed-Through Larvicides/Insect Growth Regulators (IGR)

These products can be fed to the cattle through minerals and feed and is carried through to the manure. When the horn fly eggs hatch in the manure, the larvae will be prevented from developing into adults.

These products should start to be fed 30 days before flies tend to emerge until 30 days after a killing frost. Feed-through larvicides work best if your farm is isolated from other livestock or if your neighbors have livestock they are also feeding the products. These products can impact beneficial insects, such as dung beetles.

### Fly Tags with Insecticides

Typically, ear tags provide excellent, long-term control of horn flies; some brands also reduce face fly numbers. They release small amounts of an insecticide distributed over the animal during grooming or rubbing. Be sure to pay attention to the insecticide group and do not use the same group in consecutive years. Read the label to know how many tags are needed. Often times adult cattle will require two tags while calves will require only one. Do not put the ear tags in until you can count at least 200 flies per cow. Putting them in too early will decrease their efficacy. It is also important to take the tags out after three to five months to help prevent resistance. You can often couple putting in ear tags with deworming. Most pour-on deworming products also provide external parasite control. However, you do not want to rely on dewormers as a primary external parasite control method because of the chance of the internal parasites developing resistance to the product.

Pyrethroid Group 3	Organophosphate Group 1B
Permethrin - Atroban Extra, Apollo, Deckem, Ectiban, Ear Force, Expar Extra, Gard Star Plus, New Z Permethrin, Permethrin Insecticide Ear Tags, Super Deckem II (10%) 2 tags	15% coumaphos + 35% diazinon Corathon 2 tags
10% Cyfluthrin - Cutter Gold 2 tags	Coumaphos + Diazinon - Co-Ral Plus 1 tag for horn fly, 2 tags for face fly suppression
beta-Cyfluthrin CyLence Ultra (8%) CyGuard 15% 2 tags	20% fenthion Cutter Blue 2 tags
10% z-Cypermethrin - Python, Python Magnum, ZetaGard 1 tag	pirimifos methyl Dominator 2 tags
10% l-cyhalothrin - Saber Extra/Excalibur 2 tags	New Z Diazinon (18%) Insecticide Ear Tag + Synergist
	20% diazinon OPTimizer / X-Terminator
<b>Combination Tags P + OP Groups 1B + 3</b>	40% diazinon Patriot 2 to suppress face flies
6.8 % l-Cyhalothrin + 14% Pirimiphos methyl - Double Barrel VP 2 tags	30% diazinon + 10% chlorpyrifos Warrior / Diaphos Rx 2 tags
7% Cypermethrin + 5% Chlorpyrifos - Max-Con 2 tags	
<b>Group 21</b>	<b>Abamectin Group 6</b>
15% Tolfenpyrad - Tolfenpro 1 or 2 tags	18% Abamectin XP820 2 tags

<https://entomology.ca.uky.edu/ent11>

### Dust Bags and Cattle Rubs

Dust bags and cattle rubs can be very effective for face and horn flies. The key to success with these control methods is proper placement and keeping them charged with insecticides. These devices are best placed in forced-use areas such as near mineral feeders, feed troughs, and watering sites. These can become labor-intensive as they need to be recharged with insecticide every week or so.

Fly control requires multiple approaches. These are some tools that can help control the fly population at your farm. Contact your local Extension agent if you have any questions.

## Forages for Summer Grazing

By: Paul Gonzalez, Extension Livestock Agent, N.C. Cooperative Extension, Sampson County Center

Livestock producers and horse owners who use summer annuals for grazing and/or haying should be thinking about what to plant this year. Though it may seem early, planning now will mean success later this summer. Some seed may be hard to locate, or even unavailable, which will mean choosing a second option. Most producers will probably choose the old standards, pearl millet or sorghum-sudan hybrids. Some producers are now relying on crabgrass as well. Others may even be trying "unique" forages such as lespedeza. Let's discuss each a little and you can decide what you would like to try.

Pearl millet is probably the annual of choice by most livestock producers. It is leafier than sorghum-sudan hybrids and doesn't present the risk of prussic acid poisoning. Millet grows 3 to 8 feet tall with peak production during the months of June, July, and August. It is excellent quality forage, containing 14 to 18 percentage crude protein and 60 to 65 percentage digestibility. Once allowed to head, millet is less palatable and animals consume mainly leaves. The dwarf varieties are easier to manage for grazing as there is less stem with the same number of leaves as the taller types.

The sorghum-sudan hybrids and sudangrass usually yield more than pearl millet on heavier soils. The hybrids grow 4 to 8 feet tall. Sudangrass is shorter with finer stems. Again, peak production is June, July, and August. Forage quality is good, usually 15 percent protein and about 60 percent digestibility, with high dry matter intake. Varieties out now known as brown midribbed, or BMR, seem to offer the best forage quality. The potential for prussic acid poisoning does exist from all varieties of sorghum-sudan hybrids and sudangrass. Avoid grazing young seedlings, young regrowth shoots, stunted growth, and frosted plants. There is no danger of prussic acid from feeding hay or silage from these plants.

Getting the most use from these forages can sometimes be a difficult proposition. They need to be grazed from 14 to 24 inches tall down to around 6 or 8 inches. Intensive grazing will help keep them at the proper height, as will mowing and splitting nitrogen applications. Mowing of stalks after grazing may be needed too. If cut for hay, they should be cut when growth is 30 to 40 inches tall. If bermudagrass is available, producers should consider grazing the annual and cutting the bermudagrass for hay.

More producers are testing crabgrass as summer forage. Others have made it their forage of choice. The two varieties that are available are Red River and Quick-N-Big. While it won't tolerate wet feet, crabgrass is well adapted to most soils, is drought tolerant, and responds to moisture quicker than most other summer annuals. It produces good growth from June to September with yields ranging from 3 to 5 tons per acre. It is highly palatable and usually higher quality than bermudagrass. Digestibility ranges from 62 to 72 percent and protein from 7 to 18 percent, depending on stage of growth and nitrogen fertilization.

Crabgrass can be seeded into a prepared seedbed immediately after the last spring frost. Disking or other tillage during the dormant season seems to be

necessary for productive reseeding. Nitrogen is most efficiently used in split applications at 50 to 75 pounds per acre two or three times during the growing season. An initial pre-emergence nitrogen application is very beneficial to stand development and productivity. Graze when grass is 4 inches tall. If natural reseeding is desired, the stand must be allowed to produce seed sometime during the summer. This works best with rotational grazing.

Annual lespedeza is adapted to most North Carolina soils with the exception of deep, dry sands. It tolerates low pH and fertility but will respond to fertilization. However, too much fertilizer will cause the lespedeza to be crowded out by more vigorous, higher-yielding plants in the stand. The general growth period is May to October. Peak growth occurs in June, July, and August. It is high in quality with protein levels of 14 to 18 percent and 60 to 65 percent digestibility. In fact, early cut hay may be equal to alfalfa hay. Annual lespedeza is readily grazed and has relatively high levels of phosphorus, calcium, and magnesium. Even better, it can be grazed without worrying about bloat problems. On the down side, yields are relatively low at 1.5 to 2 tons per acre in pure stands and 2 to 3 tons when mixed with grasses.

There are two main types, Kobe and Korean. Kobe type does better in the coastal plain. Both perform well in the piedmont. Marion has been the most popular variety but a new variety, called Legend, is out now that seems to be performing well in tests. In some tests, the Legend has out-yielded Marion by almost double. Ideally, seeds should be planted in February but can be sown February thru April. Seeding rate is 20 to 30 pounds per acre and should be broadcast or put out with a seeder. Drills generally put the seed too deep, as they should be no more than 1/2-inch in the ground, 1/4-inch or less being preferred. Grazing should be done in a rotational manner. If cut for hay, lespedeza should be cut in the early bloom stage. One note of caution, annual lespedeza can and will naturally reseed, so it could become somewhat of a pest under certain conditions. But don't plan on natural reseeding for the next years grazing!

One last note, don't confuse the annual lespedeza with perennial *Sericea lespedeza*. It too tolerates low fertility and acid soils but is not nearly as high in quality. It is a perennial. It will spread. It can become a nuisance. Also, if allowed to get over about 18 inches tall, it will become hard, stemmy, woody, unpalatable, quality drops rapidly, tannin levels increase, and livestock will not eat it, except goats. Goats love *Sericea*. As a bonus, the high tannin levels have been shown to work as a natural dewormer in the goats. If very tall, they will only eat the top out and you'll need to mow the tougher stems that are left. If grazed when 6 to 8 inches tall, it is eaten well by livestock. *Sericea* should be cut for hay when 12 inches tall and, whether grazed or mowed, should be taken no lower than 3 inches. Lime and fertilizer improve palatability. Seedling vigor is poor and it cannot be used the first year. *Sericea* is not typically recommended for planting but can be utilized if already on the farm.

## Are You Using the Correct Dewormer?

By: Kelly McCaskill, Extension Livestock Agent, N.C. Cooperative Extension, Moore County Center

Spring has sprung which means the dreaded “P” word is in the back of all small ruminant producers’ minds; not pollen, but parasites! If you know anything about sheep and goats, you will know that intestinal parasites are one of the biggest challenges in small ruminant production. We have some good tools in our parasite management toolbox, such as FAMACHA scoring, fecal egg counts, and conscientious grazing, as well as the use of chemical dewormers. We should be using all of the tools together in an integrated parasite management program, but often times I see producers relying primarily on dewormers to get their parasite problem under control. So how do you know which dewormer to use?

First you need to identify which type of parasite you are treating. This requires a fecal egg count. You should always work with your veterinarian when making any major herd health decisions, but depending on what county you are in, your livestock extension agent may be able to train you on how to do your own fecal egg counts. There are numerous intestinal parasites that affect small ruminants, but for deworming purposes, they can generally be clumped into three main groups: Coccidia, which is a protozoan not a worm; strongyles, aka round-worms, which includes parasites like the barber pole worm, trichostrongylus, and thread worms to name a few; and tapeworms, which are a type of flat worm. (There is a third group of true worms called flukes but they are a much less common issue in small ruminants and can be controlled by the same dewormers as tapeworms.) The fecal egg count will help you not only identify the group(s) you are battling, but help give you an idea if the use of dewormer is necessary.

Since Coccidia is a protozoan and not a true worm, there is a specific drug that is used for the treatment of coccidiosis. Corid, drug name Amprolium, is an over-the-counter treatment for coccidia, however, it is not labeled for sheep or goats, so the use of it in small ruminants requires a prescription and guidance from a veterinarian. A vet can also prescribe sulfa drugs, such as Albon or Sulmet. These drugs are all specific to only coccidia and will not treat any of the other intestinal parasites that your animals may have.

For the true worms, we use what is referred to as Anthelmintics. In small ruminants, there are three general classes of these dewormers: benzimidazoles, imidazothiazoles, and macrolides. All of these will help kill intestinal parasites but have different modes of action; think knife vs. gun vs. lead pipe; they all can do the job, but in different ways. This is an important thing to understand, because as we are seeing more and more parasites resistant to dewormers, you want to make sure you are changing classes, not just brand name or drug name, if recommended from your veterinarian or extension agent to do so.

The more commonly used benzimidazole dewormers are fenbendazole (Safeguard, Panacur) and albendazole (Valbazen); imidazothiazole dewormers are levamisole (Levisol, Tramisol) and morantel tartrate (Rumatel); and macrolide dewormers are ivermectin (Ivomec) and moxidectin (Cydectin). Of these, only fenbendazole, albendazole, and morantel tartrate are currently approved for use in goats. All others would be used as extra-label, which requires a prescription and guidance from a veterinarian.

All three classes of dewormers are labeled for some strongyles, but it varies from class to class and drug to drug as to which strongyles it will kill and how effectively it will do so. Only the benzimidazole class, or “white dewormers”, are labeled for treatment of tapeworms. *\*Note* - Valbazen has a restriction on giving to pregnant animals in the first 30 days of gestation.\*

The last part to making sure that you are giving the correct dewormer is administering the correct dosage. Since most of the dewormers that we use are not labeled for small ruminants, especially lacking for goats, you will need to consult with your veterinarian or livestock extension agent for the recommended dosage. Under-dosing is one of the biggest contributing factors to dewormer resistance, so getting your dose right is key to combating dewormer resistance on your farm.

Working closely with your veterinarian and livestock extension agent can help assure you make it through parasite season with living animals and dead worms!

## Summer Insects and Horses

By: Taylor Chavis, Extension Livestock Agent, N.C. Cooperative Extension, Robeson County Center

As we look ahead to the summer days of increasing temperature and humidity, insects start to become more prevalent and a nuisance to animals and humans alike. It's important to keep the insects under control in order to protect horses and the people on the farm.

Many of these flying insects carry diseases, so it's key for horse owners to know what bugs may be around and how to prevent them. Flies are probably the most common insect we see around our barns. Not all of them bite, but those that do can severely irritate our horses. The horses may stomp, kick, and swish their tails to get rid of the pesky bugs. Vigorous stamping can actually cause cracks in hooves from the repetitive force (according to the University of Kentucky).

There are many types of flies (house and horse are probably the ones you're familiar with) and they can all carry diseases that affect both horses and people. Stable flies have a painful bite and usually attack the flank or below the knee of animals. There is no silver bullet for fly control but reducing the number of breeding sites can significantly decrease the population. Clean barns and paddocks once a week, and spread the manure in a thin layer or compost if possible. Flies like to lay their eggs in decaying organic material, soiled bedding, and moist material, so the less opportunity you give the flies, the better your chances are. Fly traps and sticky tape are also useful to have around the barn.

Ticks are another common problem this time of year, again for both horses and humans. Ticks are often found in woody areas, brush, and overgrown grass. Eliminating these spots or restricting your animals' access to these areas can decrease their exposure. Mice are important hosts for ticks, so making sure your mice population doesn't sky rocket is key. Check the lower body, mane, and tail of your horse regularly for ticks and remove as soon as possible. If you've taken your horse out on a trail ride, definitely check your horse and yourself!

There are some less common insects you may want to consider controlling this summer. Carpenter bees do not sting often, but their loud

buzzing and aggressive behavior can certainly annoy humans and horses. They frequently make nests in eaves and window trim, so check these spots out first.

Spiders are another one we are used to seeing in horse barns but not thinking about controlling. All spiders can bite if disturbed, according to entomologist Lee Townsend from the University of Kentucky. While not all spider bites are poisonous, they do inflict a wound that can easily become infected. Of course, black widows and brown recluses are a totally different story!

Blister beetles are something we usually think about in hay. These beetles contain a chemical that is toxic to horses, and even touching them can poison your horse. Consuming too many can cause death, so this is something you need to be aware of. Oftentimes horses are exposed to these beetles when they are crushed in the hay making process; they are most often found in alfalfa hay, usually in later cuttings of hay. Your horse would have blisters in their mouth and GI tract if they consume these insects, reduced eating, and colic-like symptoms. Contact your vet immediately if your horse consumes alfalfa hay and starts exhibiting these symptoms.



Blister beetle: Photo: Clemson University/USDA Cooperative Extension

There are other insects to be concerned with on your farm, in your barns, and your pastures, but I've just highlighted a few here.

## **Youth Livestock Contests—No Critter Required!**

*By: Eve Honeycutt, Extension Livestock Agent, N.C. Cooperative Extension, Lenoir and Greene County Centers*

For youth who love livestock critters, there is no shortage of opportunities for them to show their animals. Having a calf, hog, or other project animal at home is always a learning experience. Luckily, there are plenty of other ways for critter-loving children to learn about them - and no critter is required to participate!

Contests for kids 8-18 years old:

### **4-H Skill-a-thon**

The youth livestock skill-a-thon is a contest that challenges youth individually and also as a team. The contest is divided into feed ID, breed ID, tool ID, meat cut ID, and a written test. The written test can cover anything related to livestock production, management, and the industry. Species covered are beef cattle, sheep, goats, and hogs. Participants can prepare for these contests on their own, with others, or with their Extension Agent.

### **4-H Livestock Quiz Bowl**

As with most quiz bowl contests, there are two teams of 4 members each. Questions are asked in rounds and relate to production and management of beef cattle, hogs, sheep, and goats.

### **4-H Livestock Judging**

Participants work as a team or individually on judging and selecting cattle, sheep, goats, and hogs. In addition to ranking 4 animals in a class, the youth must defend their placings by giving oral reasons to a judge. The reasons should be no longer than 2 minutes and should summarize the characteristics of the animals in the class.

For more information on 4-H livestock contests:

<https://youthlivestock.ces.ncsu.edu>

### **4-H Poultry Judging**

Poultry judging is just like livestock judging except with our feathered critters! Youth work individually and as a team to learn more about selecting live birds, as well as judging meat and eggs. Oral reasons are given for this contest as well.

### **4-H Avian Bowl**

Avian Bowl is just like our livestock quiz bowl, except the topics cover all things poultry. Teams of 4 members use a buzzer system to compete for the win.

For more information on poultry contests:

<https://poultry4hyouth.ces.ncsu.edu>

### **4-H Presentations/Congress**

A 4-H presentation is simply a way for 4-H'ers to present information they have learned. Presentations can be on any topic (including livestock and poultry) and are a great way to work on public speaking skills. There is even a category for outdoor cookery! Young grillers can delight the judges by cooking their favorite meats and telling the judges all about how to prepare them.

For more information:

<https://nc4h.ces.ncsu.edu>

## USDA Block Grant for Confined Livestock Operations

By: Richard Goforth, Specialized Poultry Agent, N.C. Cooperative Extension, South Central Area

The North Carolina Department of Agriculture and Consumer Services (NCDA&CS) is administering a disaster-related USDA Farm Service Agency grant designed for confined livestock that includes contract swine and poultry growers. The grant program will be starting soon; the fully online application opening date has not been established at the writing of this article but potential impacted growers are encouraged to start gathering information and reviewing records to determine how this grant may be able to provided them assistance. This grant program covers three different areas of loss assistance and is directly related to storm damage that occurred in 2018 from Hurricanes Florence or Michael, and weather damage from Hurricane Dorian in 2019. Growers will have to show at least a fifteen percent loss, which for confined livestock operations can be based on losses in one or more barns or houses, not just the entire operation.

One type of assistance available is for structural damages to confined livestock facilities sustained during recent weather-related disaster events. Growers of both swine and poultry who had structural damages and made repairs during 2018 or 2019 are eligible to apply for cost-share assistance of \$14 per sq. ft for poultry and \$30 per sq. ft for confined livestock, minus insurance indemnity X 50 percent of the cost.

Poultry growers that had income reductions as the result of lost growing cycles from flooding and weather-related damage can apply for an income supplement to cover part of those losses. Producers must certify that missed payment cycles were not subsidized by their integrator and provide integrator contact information for additional verification. The supplement rate is \$0.36 per bird multiplied by the number of birds per growth cycle times the number of growth cycles times 75 percent.

Another type of assistance that is available to confined livestock operations is for those that received damages to structures or losses during the covered period when part or all the facility is located within the 100-year floodplain. This cost-share grant provides funding to relocate confined livestock operations completely outside of the 100-year floodplain. The 100-year floodplain will be determined by GIS spacial view of damage due to hurricanes in 2018 or 2019. For swine operations, rebuilding remains subject to the Swine Farm Siting Act (NCGS 106-800 thru 106-806) as well as adherence to performance standards for animal waste management systems that serve swine farms. Cost-share grants are for new structures and will be calculated at \$14 per sq. ft for broiler poultry facilities and \$30 per sq. ft for swine, multiplied by 65 percent. If other assistance is received for relocation, such as from an integrator, the cost-share will be reduced.

To be eligible, growers must contribute land, capital, equipment, labor and/ or management to a farming operation or be an heir of such a producer. The attached map shows the eligible counties and year or years producers are eligible to apply for loss assistance. If a producer has eligible losses related to more than one tax identification number, they will have to fill out a separate application for each tax ID. Stay tuned for further information and alerts when the official application window opens. N.C. Cooperative Extension will be assisting with providing information to producers, as well as verifying losses in conjunction with the FSA and other agriculture agencies, to help facilitate the grant program.

